


2.2 Fuzzy Sets (Cont'd)

Crisp sets: $A = \{0, 1\}$

Concepts and thoughts are abstract and imprecise \neq random.

Fuzzy logic \rightarrow approximate knowledge



Membership function (MF) grade.

Consider:

Height:

$H = 6.001 \text{ ft}$

"Tall" MF grade: 99.9%

$H' = 5.999 \text{ ft}$

"not Tall" MF grade: 0.01%

Fuzzy set:

$$A = \{x, \mu_A(x)\}$$

$x = \text{variable } \in X$

$\mu_A = \text{MF}$

$X = \text{universe of discourse}$


1) Fuzzy sets with a discrete non-ordered universe

$$X = \{\text{Montreal}, \text{Toronto}, \text{Vancouver}\}$$

$C = \text{"desired city to live in"}$

$$C = \{(Mon, 0.5), (Tor, 0.8), (Van, 0.7)\}$$


Graphical representation:



2) Fuzzy set with a discrete ordered universe
 $X = \{0, 1, 2, 3, 4\}$

Fuzzy set A = "sensible number of cars in a family"
 $A = \{(0, 0.1), (1, 0.8), (2, 1.0), (3, 0.4), (4, 0.1)\}$


Graphical representation:



3) Fuzzy sets with a continuous space
 X = "ages" ($0 \sim 120$)

Fuzzy set B = "about 40 years old"
 $B = \{(x, \mu_B(x)), x \in X\}$

Graphical representation:



$$\mu_B(x) = \frac{1}{1 + \left(\frac{x - 40}{10}\right)^4}$$

- Subjective (X, MF)
- Not random

4) Other fuzzy set representations

For a discrete, non-ordered universe:

$$A = \frac{\mu_A(x_1)}{x_1} + \frac{\mu_A(x_2)}{x_2} + \frac{\mu_A(x_3)}{x_3} + \dots$$

e.g.

$$C = \frac{0.5}{Mon} + \frac{0.8}{Tor} + \frac{0.7}{Van}$$

For a discrete, ordered universe:

$$A = \frac{0.1}{0} + \frac{0.8}{1} + \frac{1.0}{2} + \frac{0.4}{3} + \frac{0.1}{4}$$

For a continuous space:

$$B = \frac{\mu_B(x)}{x}$$

$$B = \left[\frac{1}{1 + \left(\frac{x - 40}{10} \right)^4} \right] / x$$


If the universe space X is a continuous space, we can partition X into several fuzzy sets.

Consider:

X = "age"

Partitions:


y **M** **O**
 "young", "middle aged", "old"
 $\mu_Y(x)$, $\mu_M(x)$, $\mu_O(x)$, where $x \in X$



2.3 Other Concepts of Fuzzy Sets


1) Support

$$\text{support}(A) \rightarrow \{x | \mu_A(x) > 0\}$$



2) Core


$$\text{core}(B) \rightarrow \mu_B(x) = 1$$



3) Normality

$$\text{normality}(C) \rightarrow \max\{\mu_C(x)\} = 1$$

4) Cross-over Points



$$\mu_B(x_0) = \mu_C(x_0) = 0.5$$

x_0 = a cross-over point

this is a cross-over point,
but doesn't have specified
indication. If no specified
indication, grade = 0.5

5) Fuzzy singletons

Basically, a fuzzy set in discrete form.


Diagnosis:

Class 1, Class 2, ...



6) $\alpha - \text{cut}$

$$B = \{x, \mu_B(x) | \mu_A(x) \geq \alpha\}$$




- Strong $\alpha - \text{cut}$

$$B = \{x, \mu_B(x) | \mu_A(x) > \alpha\}$$

7) Convexity

Fuzzy sets are convex functions.




8) Fuzzy Numbers

A fuzzy number is a fuzzy set

→ normality

→ convexity (monotonically increasing, followed by monotonically decreasing, or constant)

9) Bandwidth



$$x_2 - x_1 = \text{bandwidth}$$

$$\mu_A(x_1) = \mu_A(x_2) = 0.5$$

10) Symmetry





Table 2.1: Some properties of fuzzy sets

Property name	Relation
Commutativity	$A \cap B = B \cap A$ $A \cup B = B \cup A$
Associativity	$(A \cap B) \cap C = A \cap (B \cap C)$ $(A \cup B) \cup C = A \cup (B \cup C)$
Distributivity	$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
Absorption	$A \cup (A \cap B) = A$ $A \cap (A \cup B) = A$
Idempotency (Idem = same; potent = power) (Similar to unity or identity operation)	$A \cup A = A$ $A \cap A = A$
Exclusion: Law of excluded middle Law of contradiction	$A \cup A' \subset X$ $A \cap A' \supset \emptyset$
DeMorgan's Laws	$(A \cap B)' = A' \cup B'$ $(A \cup B)' = A' \cap B'$
Boundary conditions	$A \cup X = X$ $A \cap X = A$ $A \cup \emptyset = A$ $A \cap \emptyset = \emptyset$


2.4 Set Operations

1) Subset

Consider fuzzy set A & B

If A is a subset of B :

$$A \subset B$$




2) Union (Disjunction) - OR

Given A & B

$$C = A \cup B$$

$$\mu_C(x) = \max\{\mu_A(x), \mu_B(x)\}$$




3) Intersection (Conjunction) – AND

Given A & B


$$C = A \cap B$$

$$\mu_C(x) = \min\{\mu_A(x), \mu_B(x)\}$$



4) Complement (Negation) – NOT

Given A & B



5) Cartesian Product / Co-product

$A \sim$ fuzzy set in X
 $B \sim$ fuzzy set in Y

"different universes / domains"

Cartesian product $A \times B$

is in $X \times Y$

$$\mu_{A \times B}(x, y) = \min\{\mu_A(x), \mu_B(y)\}$$


Cartesian co-product $A + B$

is in $X + Y$

$$\mu_{A+B}(x, y) = \max\{\mu_A(x), \mu_B(y)\}$$

2.4 Membership Functions (MF)

1) Triangular Membership Functions




$$\mu_A(x; a, b, c) = \begin{cases} 0 & \text{when } x \leq a \\ \frac{(x-a)}{b-a} & \text{when } a < x < b \\ \frac{(c-x)}{c-b} & \text{when } b < x < c \\ 0 & \text{when } x > c \end{cases}$$

In MATLAB:

`trimf(x, [a b c])`

2) Trapezoidal Membership Functions



$$\mu_A(x; a, b, c, d) = \begin{cases} 0 & \text{when } x < a \\ \frac{(x-a)}{b-a} & \text{when } a \leq x < b \\ 1 & \text{when } b \leq x < c \\ \frac{(d-x)}{d-c} & \text{when } c \leq x \leq d \\ 0 & \text{when } x > d \end{cases}$$


In MATLAB:

`trapmf(x, [a b c d])`

NOTE: Triangular, and trapezoidal membership functions are not continuous, which means the derivatives functions do not exist (equal to zero).

The following membership functions are continuous:

3) Gaussian Membership Functions



In MATLAB:

`gaussmf(x; [c, σ])`

$$\mu'(x) = DG(x) = e^{-\frac{1}{2}\left(\frac{x-c}{\sigma}\right)^2} \cdot \left[-\frac{1}{2} \cdot 2 \left(\frac{x-c}{\sigma} \right) \cdot \frac{1}{\sigma} \right]$$

4) Generalized Bell Membership Functions

$$\mu_A = \text{bell}(x; a, b, c) = \frac{1}{1 + \left| \frac{x-c}{a} \right|^{2b}}$$




Figure 2.8. The effects of changing parameters in bell MFs: (a) changing parameter a ; (b) changing parameter b ; (c) changing parameter c ; (d) changing a and b simultaneously but keeping their ratio constant. (MATLAB file: `allbells.m`)

In MATLAB:


`gbellmf(x; [a, b, c])`

5) Sigmoid Membership Functions

$$\mu(x) = \text{sig}(x; a, c) = \frac{1}{1 + \exp[-a(x - c)]}$$


In MATLAB:

`sigmf(x; [a, c])`



If $a > 0$, sigmf opens to the right

If $a < 0$, sigmf opens to the left



$$\mu'(x) = DS(x) = -1[1 + e^{-a(x-c)}]^{-2} e^{-a(x-c)} \cdot (-a)$$

$0 \leq x < \infty$

2.5 Fuzzy Operations

MFs $[0, 1]$

$$A \sim [0, 1]$$

$$B \sim [0, 1]$$

$$C \sim A \cup B$$

$$D \sim A \cap B$$

$$[0, 1] \times [0, 1] \rightarrow [0, 1]$$

1) Triangular Norm (T-Norm) – Generalized Intersection

$$a = \mu_A(x)$$

$$b = \mu_B(x)$$

$$T = (a, b), \quad aTb$$

Properties in table below.

2) T-Conorm (S-Norm)

Properties in table below.

Table 2.2: Some properties of a triangular norm

Item description	T-norm (triangular norm)	S-norm (T-conorm)
Function	$T: [0, 1] \times [0, 1] \rightarrow [0, 1]$	Same
Nondecreasing in each argument	If $b \geq a, d \geq c$ then $bTd \geq aTc$	Same
Commutative	$aTb = bTa$	Same
Associative	$(aTb)Tc = aT(bTc)$	Same
Boundary conditions	$aT1 = a$ $aT0 = 0$ with $a, b, c, d \in [0, 1]$	$aS0 = a$ $aS1 = 1$
Examples	Conventional: $\min(a, b)$ Product: ab Bounded max (bold intersection): $\max[0, a + b - 1]$ General: $1 - \min[1, ((1 - a)^p + (1 - b)^p)^{1/p}]$ $p \geq 1$ $\max[0, (\lambda + 1)(a + b - 1) - \lambda ab]$ $\lambda \geq -1$	Conventional: $\max(a, b)$ Set addition: $a + b - ab$ Bounded min (bold union): $\min[1, a + b]$ General: $\min(1, (a^p + b^p)^{1/p})$ $p \geq 1$ $\min[1, a + b + \lambda ab]$ $\lambda \geq -1$
DeMorgan's Laws	$aSb = 1 - (1 - a) T(1 - b)$ $aTb = 1 - (1 - a) S(1 - b)$	

Example 2.3 (Similar to Example 2.13)

Use DeMorgan's law to determine the S-norm corresponding to $\max(x, y)$, and T-norm corresponding to $\min(x, y)$.

Solution 2.3

$$\begin{aligned}xSy &= 1 - (1 - x)T(1 - y) \\T \rightarrow \min & \\&= 1 - \min[(1 - x), (1 - y)] \\&= \begin{cases} 1 - (1 - y) = y; & x < y \\ 1 - (1 - x) = x; & x \geq y \end{cases} \\xSy &= \max(x, y)\end{aligned}$$

Example 2.4 (Similar to Example 2.14)

Prove that the min operator is the largest T-norm and the max operator is the smallest S-norm.

Solution 2.4

Nondecreasing, boundary conditions

$$\begin{aligned}xTy &\leq 1Ty = y \\xTy &\leq xT1 = x \\xTy &\leq \min(x, y)\end{aligned}$$