

Sept. 14 / 16

P - Rank of a coefficient matrix.

Q - Rank of an augmented matrix

N - Number of unknowns

M - Number of equations

 $N = M ?$ When $P = Q = N$ a system has a unique solution $P < Q$ a system has no solution
 $\begin{cases} N > P = Q \\ S = N - P \end{cases}$ a system has infinitely many solutions
 number of free variables.

e.g.

$$\left(\begin{array}{ccc|c} 1 & 0 & 0 & a \\ 0 & 1 & 0 & b \\ 0 & 0 & 1 & c \end{array} \right) \quad P = Q = N$$

$$\left(\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right) \quad P < Q$$

For example:

$$\left(\begin{array}{ccc|c} 1 & 2 & -4 & 6 \\ 0 & 1 & 2 & 5 \\ 0 & 0 & 1 & 3 \end{array} \right) \quad \begin{aligned} x + 2y - 4z &= 6 & (\text{eq. 1}) \\ + 1y + 2z &= 5 & (\text{eq. 2}) \\ + z &= 3 & (\text{eq. 3}) \end{aligned}$$

(the augmented matrix is in the echelon form)

 $P = Q = N \therefore$ System has unique solution.

Solution:

- Substitute (eq. 3) into (eq. 2) to solve for "y"
- Substitute "y" and "z" into (eq. 1) to solve for "x"

(2)

Sept. 14/16

For a matrix, the first number represents rows.
 $(m \times n)$ -matrix A = m-rows, n-columns

Sum of Matrices

$$A + B = \begin{pmatrix} a_{11} & \dots & a_{1n} \\ \vdots & & \vdots \\ a_{m1}, & \dots & a_{mn} \end{pmatrix} + \begin{pmatrix} b_{11} & \dots & b_{1n} \\ \vdots & & \vdots \\ b_{m1}, & \dots & b_{mn} \end{pmatrix}$$

(should be)

$$cA = \begin{pmatrix} ca_{11} & \dots & ca_{1n} \\ \vdots & & \vdots \\ ca_{m1}, & \dots & ca_{mn} \end{pmatrix}$$

(2)

$$\text{Let } A = \begin{pmatrix} -1 & 2 \\ 2 & -3 \\ 0 & 1 \\ 4 & -1 \end{pmatrix}_{4 \times 2} \quad \text{Let } B = \begin{pmatrix} 10 & -1 & 0 \\ 1 & 2 & 1 \end{pmatrix}_{2 \times 3}$$

Sept. 16 / 16

Find AB .

$$AB = (m \times r)(r \times n) = m \times n$$

$$= (4 \times 2)(2 \times 3) = (4 \times 3)$$

COLUMN 1

$$\begin{aligned} \text{so..} &= (-1)(10) + (2)(1) \Rightarrow -8 \\ &= (2)(10) + (-3)(1) \Rightarrow 17 \\ &= (0)(10) + (1)(1) \Rightarrow 1 \\ &= (4)(10) + (-1)(1) \Rightarrow 39 \end{aligned}$$

①

Sept. 16 / 16

Determine which matrices are in reduced echelon form and which are only in echelon form.

$$\left(\begin{array}{cccc} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right) \text{ reduced echelon form}$$

$$\left(\begin{array}{ccc} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{array} \right) \text{ reduced echelon form}$$

$$\left(\begin{array}{cc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{array} \right) \text{ not echelon form}$$

$$\left(\begin{array}{ccccc} 1 & 0 & 1 & 1 & 1 \\ 0 & 2 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 4 \\ 0 & 0 & 0 & 0 & 5 \end{array} \right) \text{ echelon form}$$

$$\begin{aligned} x_1 - 2x_2 &= -1 \\ 3x_1 + 5x_2 &= 30 \\ 5x_1 + x_2 &= 28 \end{aligned}$$

$$\left(\begin{array}{cc|c} 1 & -2 & -1 \\ 3 & 5 & 30 \\ 5 & 1 & 28 \end{array} \right) \xrightarrow{\begin{array}{l} -3R_1 + R_2 \\ -5R_1 + R_3 \end{array}}$$

Augmented Matrix

$$\left(\begin{array}{cc|c} 1 & -2 & -1 \\ 0 & 11 & 33 \\ 0 & 11 & 33 \end{array} \right) \xrightarrow{-R_2 + R_3} \left(\begin{array}{cc|c} 1 & -2 & -1 \\ 0 & 11 & 33 \\ 0 & 0 & 0 \end{array} \right)$$

echelon form

(2)

$$x_1 - 2x_2 = -1$$

$$11x_2 = 33$$

$$x_2 = 3$$

$$x_1 - 2x_2 = -1$$

$$x_1 = -1 + 2(3)$$

$$x_1 = 5$$

∴ the system has a unique solution ($P = q = n$)

$$\begin{matrix} P = 2 \\ q = 2 \\ n = 2 \end{matrix}$$

Example 2

$$6x_1 - 5x_2 - 3x_3 = 55$$

$$x_1 - x_2 - x_3 = 8$$

$$7x_1 - 6x_2 - 4x_3 = 63$$

$$\left(\begin{array}{ccc|c} 6 & -5 & -3 & 55 \\ 1 & -1 & -1 & 8 \\ 7 & -6 & -4 & 63 \end{array} \right) \xrightarrow{\text{reorganize}} \left(\begin{array}{ccc|c} 1 & -1 & -1 & 8 \\ 6 & -5 & -3 & 55 \\ 7 & -6 & -4 & 63 \end{array} \right) \dots$$

$$\xrightarrow{\begin{matrix} -6R_1 + R_2 \\ -7R_1 + R_3 \end{matrix}} \left(\begin{array}{ccc|c} 1 & -1 & -1 & 8 \\ 0 & 1 & 3 & 7 \\ 0 & 1 & 3 & 7 \end{array} \right) \xrightarrow{-R_2 + R_3} \left(\begin{array}{ccc|c} 1 & -1 & -1 & 8 \\ 0 & 1 & 3 & 7 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

$$\xrightarrow{\text{echelon form}} \left(\begin{array}{ccc|c} 1 & -1 & -1 & 8 \\ 0 & 1 & 3 & 7 \\ 0 & 0 & 0 & 0 \end{array} \right) \quad \begin{matrix} P = 2 \\ q = 2 \\ n = 3 \end{matrix}$$

echelon form

~~UNIQUE~~

$$x_1 - x_2 - x_3 = 8 \quad = \quad x_1 - (7 - 3x_3) - x_3 = 8$$

$$x_2 + 3x_3 = 7 \quad \Rightarrow \quad x_2 = 7 - 3x_3$$

... where x_3 is any real number.

$$x_1 = 7 - 3x_3$$

$$x_2 = 7 - 3x_3$$

infinitely many solutions.

(3)

$$\begin{array}{l} x_1 + 3x_2 + x_3 = 0 \\ 4x_1 + 2x_2 - x_3 = 0 \\ -x_1 + x_2 + x_3 = 1 \end{array} \Rightarrow \left(\begin{array}{ccc|c} 1 & 3 & 1 & 0 \\ 4 & 2 & -1 & 0 \\ -1 & 1 & 1 & 1 \end{array} \right) \dots$$

$$\dots \xrightarrow{\frac{-4R_1 + R_2}{R_1 + R_3}} \left(\begin{array}{ccc|c} 1 & 3 & 1 & 0 \\ 0 & -10 & -5 & 0 \\ 0 & 4 & 2 & 1 \end{array} \right) \xrightarrow{(\frac{1}{5})R_2} \left(\begin{array}{ccc|c} 1 & 3 & 1 & 0 \\ 0 & 2 & 1 & 0 \\ 0 & 4 & 2 & 1 \end{array} \right) \dots$$

$$\dots \xrightarrow{-2R_2 + R_3} \left(\begin{array}{ccc|c} 1 & 3 & 1 & 0 \\ 0 & 2 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right) \quad P=2 \quad Q=3$$

$$0x_1 + 0x_2 + 0x_3 = 1$$

∴ the system has no solution.

Compute the Matrix: $3A + 4B$ where $A = \begin{pmatrix} 3 & -5 \\ 2 & 7 \end{pmatrix}$

$$\text{and } B = \begin{pmatrix} -1 & 0 \\ 3 & -4 \end{pmatrix}$$

$$3A = \begin{pmatrix} 9 & -15 \\ 6 & 21 \end{pmatrix}_{2 \times 2} \quad 4B = \begin{pmatrix} -4 & 0 \\ 12 & -16 \end{pmatrix}_{2 \times 2}$$

$$\begin{aligned} 3A + 4B &= \begin{pmatrix} 9 & -15 \\ 6 & 21 \end{pmatrix} + \begin{pmatrix} -4 & 0 \\ 12 & -16 \end{pmatrix} \Rightarrow \begin{pmatrix} 9-4 & -15-0 \\ 6+12 & 21+(-16) \end{pmatrix} \\ &= \begin{pmatrix} 5 & -15 \\ 18 & 5 \end{pmatrix} \end{aligned}$$

(4)

Calculate AB and BA

$$A = (1 \ 2 \ 3)_{1 \times 3}$$

$$B = \begin{pmatrix} 3 \\ 4 \\ 5 \end{pmatrix}_{3 \times 1}$$

 AB

$$(1 \times 3)(3 \times 1) = (1 \times 1)$$

$$= (1 \ 2 \ 3) \begin{pmatrix} 3 \\ 4 \\ 5 \end{pmatrix} = (1(3) + 2(4) + 3(5)) = (26)_{1 \times 1}$$

 BA

$$(3 \times 1)(1 \times 3) = (3 \times 3)$$

$$= \begin{pmatrix} 3 \\ 4 \\ 5 \end{pmatrix} (1 \ 2 \ 3) = \begin{pmatrix} 3(1) & 3(2) & 3(3) \\ 4(1) & 4(2) & 4(3) \\ 5(1) & 5(2) & 5(3) \end{pmatrix}$$

$$= \begin{pmatrix} 3 & 6 & 9 \\ 4 & 8 & 12 \\ 5 & 10 & 15 \end{pmatrix}_{3 \times 3}$$

$$A = \begin{pmatrix} 8 \\ -2 \end{pmatrix}_{2 \times 1}$$

$$B = \begin{pmatrix} 0 & -2 \\ 3 & 1 \\ -4 & 5 \end{pmatrix}_{3 \times 2}$$

(row \times column)For AB

$$(2 \times 1)(3 \times 2)$$

~~The product is not defined.~~

(5)

BA

$$(3 \times 2) (2 \times 1) = (3 \times 1) \Rightarrow \begin{pmatrix} 0 & -2 \\ 3 & 1 \\ -4 & 5 \end{pmatrix} \begin{pmatrix} 3 \\ -2 \end{pmatrix} \Rightarrow$$

$$\begin{pmatrix} 0(3) + (-2)(-2) \\ 3(3) + 1(-2) \\ -4(3) + 5(-2) \end{pmatrix} \Rightarrow \begin{pmatrix} 0 + (+4) \\ 9 + (-2) \\ -12 + (-10) \end{pmatrix} \Rightarrow \begin{pmatrix} 4 \\ 7 \\ -22 \end{pmatrix}_{3 \times 1}$$

Identity Matrix $I_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$

$$AI = \begin{pmatrix} 2 & 3 & 4 \\ 5 & 6 & 7 \\ 8 & 9 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$IA = A$$

Example:

$$-x_1 - 2x_2 + x_3 = 5$$

$$AX = B$$

$$-2x_1 + 3x_2 + x_3 = 1$$

A - coefficient matrix

$$-x_1 + 3x_2 + 2x_3 = 2$$

X - column vector of unknowns

B - column vector of constants

$$\left(\begin{array}{ccc|c} -1 & -2 & 1 & 5 \\ -2 & 3 & 1 & 1 \\ -1 & 3 & 2 & 2 \end{array} \right) \left(\begin{array}{c} x_1 \\ x_2 \\ x_3 \end{array} \right)$$

A — B — X

$$\left(\begin{array}{ccc|c} -1 & -2 & 1 & 5 \\ -2 & 3 & 1 & 1 \\ -1 & 3 & 2 & 2 \end{array} \right) \rightarrow ? \text{ answer on board resolves...}$$

$$\left(\begin{array}{ccc|c} 1 & -2 & 1 & 5 \\ -2 & 3 & 1 & 1 \\ 1 & 3 & 2 & 2 \end{array} \right) \text{ diff. matrix?}$$

(6)

$$x + y - 2z + 4w = 5$$

$$2x + 2y - 3z + w = 3$$

$$3x + 3y - 4z - 2w = 1$$

Ax

$$\left(\begin{array}{cccc|c} 1 & 1 & -2 & 4 & 5 \\ 2 & 2 & -3 & 1 & 3 \\ 3 & 3 & -4 & -2 & 1 \end{array} \right) \xrightarrow{\text{A } 3 \times 4 \quad 3 \times 1} \left(\begin{array}{c} x \\ y \\ z \\ w \end{array} \right) \quad x \quad 4 \times 1$$

$$\therefore Ax = 3 \times 1$$

$$\left(\begin{array}{cccc|c} 1 & 1 & -2 & 4 & 5 \\ 2 & 2 & -3 & 1 & 3 \\ 3 & 3 & -4 & -2 & 1 \end{array} \right) \xrightarrow{\begin{array}{l} -2R_1 + R_2 \\ -3R_1 + R_3 \end{array}} \left(\begin{array}{cccc|c} 1 & 1 & -2 & 4 & 5 \\ 0 & 0 & 1 & -7 & -7 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right)$$

$$x + y - 2z + 4w = 5$$

$$z - 7w = -7$$

$$\left(\begin{array}{c} x \\ y \\ z \\ w \end{array} \right) = \left(\begin{array}{cccc} -9 & -y & +10w \\ 0 & 5 & +7w \\ -7 & 0 & 0 \\ 0 & 0 & w \end{array} \right)$$

$$z = -7 + 7w$$

$$x + y - 2(-7 + 7w) + 4w = 5$$

$$x = -y + 2(-7 + 7w) - 4w + 5$$

$$x = -y - 14 + 14w - 4w + 5$$

$$x = -9 - y + 10w$$

$$\left(\begin{array}{c} -9 \\ 0 \\ -7 \\ 0 \end{array} \right) + \left(\begin{array}{c} -y \\ 5 \\ 0 \\ 0 \end{array} \right) + \left(\begin{array}{c} 10w \\ 0 \\ 7w \\ w \end{array} \right)$$

$$x = x_0 + yx_1 + wx_2$$

$$\left(\begin{array}{c} x \\ y \\ z \\ w \end{array} \right) = \left(\begin{array}{c} -9 \\ 0 \\ -7 \\ 0 \end{array} \right) + y \left(\begin{array}{c} -1 \\ 1 \\ 0 \\ 0 \end{array} \right) + w \left(\begin{array}{c} 10 \\ 0 \\ 7 \\ 1 \end{array} \right)$$

$$x = x_0 + yx_1 + wx_2$$

where ~~x_0~~ , y, w

are any real number.