

4.2.2 Transform OF Derivatives

Nov. 13 / 17

Applied Anal.

$$\mathcal{L}\{f'(t)\} = -f'(0) + s \mathcal{L}\{f(t)\}$$

$$\begin{aligned}\mathcal{L}\{f''(t)\} &= \mathcal{L}\{f'(t)'\} \\ &= -f'(0) + s \mathcal{L}\{f'(t)\} \\ &= -f'(0) + s[-f(0) + s \mathcal{L}\{f(t)\}] \\ &= s^2 \mathcal{L}\{f(t)\} - sf(0) - f'(0)\end{aligned}$$

$$\mathcal{L}\{f'''(t)\} = s^3 \mathcal{L}\{f(t)\} - s^2 f(0) - sf'(0) - f''(0)$$

Thm. 4.2.2 If $f, f', \dots, f^{(n-1)}$ are continuous on $[0, +\infty)$ and of exponential order and if $f^{(n)}$ is piecewise continuous on $[0, \infty)$ - then;

$$\mathcal{L}\{f^{(n)}(t)\} = s^n F(s) - s^{n-1} f(0) - s^{n-2} f'(0) - \dots - f^{(n-1)}(0)$$

$$\text{where } F(s) = \mathcal{L}\{f(t)\}$$

Solve the linear DE's

→ Ex. Use the laplace transform to solve
 $\frac{dy}{dt} - 2y = 4 \cos 2t, \quad y(0) = 2$

$$\text{Solution} \quad \mathcal{L}\{y' - 2y\} = \mathcal{L}\{4 \cos 2t\}$$

$$\mathcal{L}\{y'\} - 2 \mathcal{L}\{y\} = 4 \mathcal{L}\{\cos 2t\}$$

$$\left[-y(0) + s \mathcal{L}\{y\} \right] - 2 \mathcal{L}\{y\} = 4 \cdot \frac{s}{s^2 + 4}$$

$$-2 + sY(s) - 2Y(s) = \frac{4s}{s^2 + 4}$$

$$sY(s) - 2Y(s) = 2 + \frac{4s}{s^2 + 4}$$

$$Y(s)(s-2) = 2 + \frac{4s}{s^2 + 4} = \frac{2(s^2 + 4) + 4s}{s^2 + 4}$$

$$Y(s) = \frac{2s^2 + 4s + 8}{(s-2)(s^2 + 4)}$$

$$Y = \mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{2s^2 + 4s + 8}{(s-2)(s^2 + 4)}\right\}$$

$$\frac{2s^2 + 4s + 8}{(s-2)(s^2 + 4)} = \frac{A}{(s-2)} + \frac{Bs + C}{(s^2 + 4)}$$

$$2s^2 + 4s + 8 = A(s^2 + 4) + (Bs + C)(s - 2)$$

$$(1) s=2; \quad 8 + 8 + 8 = 8A + 0; \quad A = 3$$

$$(2) \text{coeff. for } s^2; \quad 2 = A + B; \quad B = -1$$

$$(3) \text{constant term}; \quad 8 = 4A - 2C; \quad C = 2$$

$$\begin{aligned} \mathcal{L}\{Y(s)\} &= \mathcal{L}^{-1}\left\{\frac{3}{s-2} + \frac{-s+2}{s^2+4}\right\} \\ &= 3\mathcal{L}^{-1}\left\{\frac{1}{s-2}\right\} + -1\mathcal{L}^{-1}\left\{\frac{s}{s^2+4}\right\} + \mathcal{L}^{-1}\left\{\frac{2}{s^2+4}\right\} \\ &= 3e^{2t} - \cos 2t + \sin 2t \end{aligned}$$

Ex. Solve the IVP Laplace Transform

$$y'' + 5y' - 14y = e^{-3t}, \quad y(0) = 1, \quad y'(0) = 2$$

$$\underline{\text{Solution}} \quad \mathcal{L}\{y''\} + 5\mathcal{L}\{y'\} - 14\mathcal{L}\{y\} = \mathcal{L}\{e^{-3t}\}$$

$$[s^2 Y(s) - s y(0) - y'(0)] + 5[s y(s) - y(0)] - 14 Y(s) = \frac{1}{s+3}$$

$$s^2 Y(s) - s - 2 + 5s Y(s) - 5 - Y(s) = \frac{1}{s+3}$$

$$s^2 Y(s) + 5s Y(s) - 14 Y(s) = s + 7 + \frac{1}{s+3}$$

$$Y(s) (s^2 + 5s - 14) = s + 7 + \frac{1}{s+3} = \frac{(s+7)(s+3)+1}{s+3}$$

$$Y(s) = \frac{s^2 + 10s + 22}{(s+3)(s^2 + 5s - 14)}$$

$$y(t) = \mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{s^2 + 10s + 22}{(s+3)(s^2 + 5s - 14)}\right\}$$

$$\frac{s^2 + 10s + 22}{(s+3)(s-2)(s+7)} = \frac{A}{(s+3)} + \frac{B}{(s-2)} + \frac{C}{(s+7)}$$

$$s^2 + 10s + 22 = A(s-2)(s+7) + B(s+3)(s+7) + C(s+3)(s-2)$$

$$- \text{ for } s=2; \quad 4 + 20 + 22 = 45B; \quad B = 46/45$$

$$- \text{ for } s=-3; \quad 9 - 30 + 22 = -20A; \quad A = -1/20$$

$$- \text{ for } s=-7; \quad 49 - 70 + 22 = 36C; \quad C = 1/36$$

$$\begin{aligned} y(t) &= \mathcal{L}^{-1}\left\{-\frac{1}{20} \frac{1}{s+3}\right\} + \mathcal{L}^{-1}\left\{\frac{46}{45} \frac{1}{s-2}\right\} + \mathcal{L}^{-1}\left\{\frac{1}{36} + \frac{1}{s+7}\right\} \\ &= -\frac{1}{20} e^{-3t} + \frac{46}{45} e^{2t} + \frac{1}{36} e^{-7t} \end{aligned}$$

Thm. 4.2.3 (Behavior of $F(s)$ as $s \rightarrow \infty$)

If F is piecewise continuous on $[0, \infty)$

and of exponential order, then $\lim_{s \rightarrow \infty} \mathcal{L}\{f(t)\} = 0$
 i.e. If $\mathcal{L}\{F(s)\}$ exist, then

$$\lim_{s \rightarrow \infty} F(s) = 0$$

Example: Let $F(s) = \frac{s}{s+1}$: then,

$$\lim_{s \rightarrow \infty} \frac{s}{s+1} = 1 \neq 0. \quad \text{So } \mathcal{L}\{F(s)\} \text{ DNE}$$

4.3 Translation Theorems

Nov. 15 / 17
Applied Anal.

4.3.1 Translation on the s-axis

Thm 4.3.1 (First Translation Theorem)

If $\mathcal{L}\{f(t)\} = F(s)$, and a is any real number,
then; $\mathcal{L}\{e^{at} f(t)\} = F(s-a)$

Proof: $\mathcal{L}\{e^{at} f(t)\} = \int_0^\infty e^{-st} [e^{at} f(t)] dt$
 $\Rightarrow \int_0^\infty e^{-(s-a)t} f(t) dt = F(s-a)$

Notation: $F(s-a) = F(s) \Big|_{s \rightarrow s-a}$
 $\mathcal{L}\{e^{at} f(t)\} = \mathcal{L}\{f(t)\} \Big|_{s \rightarrow s-a}$

→ Ex. Use the first translation theorem to evaluate: $\mathcal{L}\{e^{2t} \cdot t^4\}$

$$\Rightarrow \mathcal{L}\{t^4\} \Big|_{s \rightarrow s-2}$$

$$\Rightarrow \frac{4!}{s^5} \Big|_{s \rightarrow s-2} \Rightarrow \frac{4 \cdot 3 \cdot 2 \cdot 1}{s^5} \Big|_{s \rightarrow s-2} \Rightarrow \frac{24}{(s-2)^5}$$

→ Ex: $\mathcal{L}\{e^{-3t} \sin 4t\} = \mathcal{L}\{\sin 4t\} \Big|_{s \rightarrow s-(-3)}$

$$\Rightarrow \frac{4}{s^2 + 16} \Big|_{s \rightarrow s+3} \Rightarrow \frac{4}{(s+3)^2 + 16}$$

$$\mathcal{L}^{-1}\{F(s-a)\} = \mathcal{L}^{-1}\{F(s) \Big|_{s \rightarrow s-a}\}$$

$$\mathcal{L}^{-1}\{F(s) \Big|_{s \rightarrow s-a}\} = e^{at} \mathcal{L}^{-1}\{F(s)\}$$

→ Ex. $\mathcal{L}^{-1}\left\{\frac{1}{(s-2)^2}\right\} = \mathcal{L}^{-1}\left\{\frac{1}{s^2} \Big|_{s \rightarrow s-2}\right\}$
 $= e^{2t} \mathcal{L}^{-1}\left\{\frac{1}{s^2}\right\} = e^{2t} \cdot t$

→ Ex. $\mathcal{L}^{-1}\left\{\frac{(2s+3)}{(s-5)^2}\right\}$ $\tilde{s} = s-5$ $s = \tilde{s} + 5$

$\frac{2s+3}{s^2} = \frac{2(\tilde{s}+5)+3}{\tilde{s}^2}$

$$\begin{aligned}
 &= \mathcal{L}^{-1} \left\{ \frac{2(s-5) + 13}{(s-5)^2} \right\} \\
 &= \mathcal{L}^{-1} \left\{ \frac{2s+13}{s^2} \mid s \rightarrow s-5 \right\} \\
 \Rightarrow e^{st} \cdot \mathcal{L}^{-1} \left\{ \frac{2s+13}{s^2} \right\} &= e^{st} \mathcal{L}^{-1} \left\{ \frac{2s}{s^2} \right\} + e^{st} \mathcal{L}^{-1} \left\{ \frac{13}{s^2} \right\} \\
 &\Rightarrow e^{st} (2 + 13t)
 \end{aligned}$$

$$\begin{aligned}
 \rightarrow \text{Ex: } \mathcal{L}^{-1} \left\{ \frac{2s+7}{s^2+6s+9} \right\} &= \mathcal{L}^{-1} \left\{ \frac{2s+7}{(s+3)^2} \right\} \\
 \Rightarrow \frac{2s+7}{(s+3)(s+4)} &= \frac{A}{(s+3)} + \frac{B}{(s+4)} \quad \dots
 \end{aligned}$$

$$\begin{array}{l}
 \mathcal{L}^{-1} \left\{ \frac{2s+7}{s^2+6s+10} \right\} \\
 \mathcal{L}^{-1} \left\{ \frac{2s+7}{(s+3)^2+1} \right\} \\
 \mathcal{L}^{-1} \left\{ \frac{2(s-3)+7}{s^2+1} \right\} \\
 \Rightarrow \mathcal{L}^{-1} \left\{ \frac{2s+1}{s^2+1} \right\} \\
 \Rightarrow \mathcal{L}^{-1} \left\{ \frac{2s+1}{s^2+1} \mid s \rightarrow s+3 \right\} \\
 \Rightarrow e^{-3t} \mathcal{L}^{-1} \left\{ \frac{2s+1}{s^2+1} \right\} = e^{-3t} \left[\mathcal{L}^{-1} \left\{ \frac{2s}{s^2+1} \right\} + \mathcal{L}^{-1} \left\{ \frac{1}{s^2+1} \right\} \right] \\
 \Rightarrow e^{-3t} [2 \cos t + \sin t]
 \end{array}$$

\rightarrow Ex. Solve the IVP

$$y'' - 4y' + 4y = t^2 e^{2t}, \quad y(0) = 1, \quad y'(0) = 1$$

$$\text{Solution} \quad \mathcal{L}\{y''\} - 4\mathcal{L}\{y'\} + 4\mathcal{L}\{y\} = \mathcal{L}\{t^2 e^{2t}\}$$

$$[s^2 Y(s) - s y(0) - y'(0)] - 4[s Y(s) - y(0)] + 4 Y(s) \dots$$

$$\dots = \mathcal{L}\{t^2 \mid s \rightarrow s-2\}$$

$$s^2 Y(s) - s - 1 - 4(s Y(s) - 1) + 4(Y(s)) = \frac{2!}{s^{2+2}} \mid s \rightarrow s-2$$

$$s^2 Y(s) - s - 1 - 4s Y(s) + 4 + 4 Y(s) = \frac{2}{(s-2)^3}$$

$$Y(s^2 - 4s + 4) - s + 3 = \frac{2}{(s-2)^3}$$

$$\begin{aligned}
 Y(s) &= \frac{s-3}{(s^2-4s+4)} + \frac{1}{(s^2-4s+4)(s-2)^3} \\
 Y(t) &= \mathcal{L}^{-1} \left\{ \frac{s-3}{(s-2)^2} + \frac{1}{(s-2)^3} \right\} \\
 &= \mathcal{L}^{-1} \left\{ \frac{(s-2)-1}{(s-2)^2} + \frac{1}{(s-2)^3} \right\} \\
 &= \mathcal{L}^{-1} \left\{ \frac{s-1}{(s-2)^2} + \frac{1}{s^3} \Big| s \rightarrow s-2 \right\} \\
 &= e^{2t} \mathcal{L}^{-1} \left\{ \frac{s-1}{s^2} + \frac{1}{s^3} \right\} \\
 &= e^{2t} \left[\mathcal{L}^{-1} \left\{ \frac{1}{s} - \frac{1}{s^2} \right\} + \mathcal{L}^{-1} \left\{ \frac{1}{s^3} \right\} \right] \\
 &= e^{2t} \left[1 - t + \mathcal{L}^{-1} \left\{ \frac{4!}{s^4} \cdot \frac{1}{4!} \right\} \right] \\
 &= e^{2t} \left(1 - t + \frac{1}{4!} \cdot t^4 \right)
 \end{aligned}$$

Ex. Solve $y'' + 4y' + 5y = 3 + e^{-2t}$

$$y(0) = 0, \quad y'(0) = 0$$

$$\begin{aligned}
 \text{Solution} \quad & \mathcal{L} \{ y'' \} + 4 \mathcal{L} \{ y' \} + 5 \mathcal{L} \{ y \} = \mathcal{L} \{ 3 + e^{-2t} \} \\
 (5^2 Y(s) - 5Y(0) - Y'(0)) + 4(5Y(s) - Y'(0)) + 5(Y(s)) &= \frac{3}{s} + \frac{1}{s+2} \\
 5^2 (Y(s)) + 4s(Y(s)) + 5(Y(s)) &= \frac{3}{s} + \frac{1}{s+2} \\
 Y(s) (s^2 + 4s + 5) &= \frac{3}{s} + \frac{1}{s+2} = \frac{3(s+2) + 5}{s(s+2)}
 \end{aligned}$$


$$Y(s) = \frac{4s+6}{s(s+2)(s^2+4s+5)}$$

4.3.2 - Translation on the t -axis

$f(t-a)$?

Def. The Unit step Function

$$u(t-a) = \begin{cases} 0 & \text{if } 0 \leq t < a \\ 1 & \text{if } t \geq a \end{cases}$$



Ex. $f(t) = \begin{cases} 0 & \text{if } 0 \leq t < 3 \\ \frac{1}{2}t & \text{if } t \geq 3 \end{cases} = \frac{1}{2}t \cdot u(t-3)$

$$1 - u(t-a) = \begin{cases} 1 & \text{if } 0 \leq t < a \\ 0 & \text{if } t \geq a \end{cases}$$

Ex. $g(t) = \begin{cases} t^2, & 0 \leq t < 3 \\ 0, & t \geq 3 \end{cases} = t^2 \cdot (1 - u(t-3))$


Ex. $h(t) = \begin{cases} t^2, & 0 \leq t < 3 \\ \frac{1}{2}t, & t \geq 3 \end{cases}$

$$\Rightarrow \begin{cases} t^2, & 0 \leq t < 3 \\ 0, & t \geq 3 \end{cases} + \begin{cases} 0 & 0 \leq t < 3 \\ \frac{1}{2}t & t \geq 3 \end{cases}$$

$$= t^2(1 - u(t-3)) + \frac{1}{2}t \cdot u(t-3)$$

If $a < b$

$$u(t-a) - u(t-b) = \begin{cases} 0, & 0 \leq t < a \\ 1, & a \leq t < b \\ 0, & t \geq b \end{cases}$$



$$\text{Ex. } f(t) = \begin{cases} 0, & 0 \leq t < 1 \\ e^t, & 1 \leq t < 3 \\ 0, & t \geq 3 \end{cases}$$

$$= e^t [u(t-1) - u(t-3)]$$

$$f(t-a)u(t-a) = \begin{cases} 0, & 0 \leq t < a \\ f(t-a), & t \geq a \end{cases}$$

Thm. 4.3.2 (Second Translation Theorem)

$$\text{For } a > 0, \quad \mathcal{L}\{f(t-a)u(t-a)\} = e^{-as} \mathcal{L}\{f(t)\}$$

$$\text{or } \mathcal{L}\{f(t)u(t-a)\} = e^{as} \mathcal{L}\{f(t-a)\}$$

$$\mathcal{L}\{f(t+a-a)\}u(t-a) = e^{-as} \mathcal{L}\{f(t+a)\}$$

$$\begin{aligned} \text{PROOF: } \mathcal{L}\{f(t-a)u(t-a)\} &= \int_0^\infty e^{-st} f(t-a)u(t-a) dt \\ &= \int_0^a e^{-st} f(t-a)u(t-a) dt + \int_a^\infty e^{-st} f(t-a)u(t-a) dt \\ &= \int_a^\infty e^{-st} f(t-a) dt \underset{u=t-a}{=} \int_0^\infty e^{-s(u+a)} f(u) du \\ &= \int_0^\infty e^{-su} \cdot e^{-sa} f(u) du \\ &= e^{-as} \int_0^\infty e^{-su} f(u) du = e^{-as} \mathcal{L}\{f(t)\} \end{aligned}$$

$$\boxed{\mathcal{L}\{f(t-a)u(t-a)\} = e^{-as} \mathcal{L}\{f(t)\}}$$

$$\boxed{\mathcal{L}^{-1}\{e^{-as} F(s)\} = f(t-a)u(t-a)}$$

$$\text{where } f(t) = \mathcal{L}^{-1}\{F(s)\}$$

$$\text{Ex. } \mathcal{L}\{u(t-2)\} = e^{-2s} \mathcal{L}\{1\} = \frac{e^{-2s}}{s}$$

$$\text{Ex. } \mathcal{L}^{-1}\left\{\frac{1}{s-2} e^{-3s}\right\} = f(t-3)u(t-3)$$

$$\text{where } f(t) = \mathcal{L}^{-1}\left\{\frac{1}{s-2}\right\} = e^{2t}$$

$$\mathcal{L}^{-1}\left\{\frac{1}{s-2} e^{-3s}\right\} = e^{2(t-3)}u(t-3) = e^{2t-6}u(t-3)$$

$$= \begin{cases} 0, & 0 \leq t < 3 \\ e^{2s-6}, & t \geq 3 \end{cases}$$